
120. Shoulder from behind. The Edinburgh Stereoscopic Atlas of Anatomy. Upper Limb. Surface anatomy - N°3. Upper part of the back.

Numéro d'inventaire : 1979.26251.1

Type de document : photographie

Éditeur : Edinburgh Stereoscopic Anatomy, T.C. & E.C. Jack (34, Henrietta St., London, W.C. Edinburgh, London)

Période de création : 1er quart 20e siècle

Date de création : 1900 (vers)

Description : Epreuves stéréoscopiques positives gélatino-argentiques contrecollées sur un carton rigide contenant un texte descriptif (format du carton : 230 x 180). Série rangée dans un emboîtement en carton sous forme de reliure en deux parties avec la mention "Pestalozzi Stereographs. Anatomy" sur la tranche.

Mesures : hauteur : 90 mm ; largeur : 180 mm

Notes : Descriptif : la partie supérieure du dos (anatomie).

Mots-clés : Méthodes pédagogiques actives (y compris la coopération scolaire, classes vertes, méthode Freinet)

Pratique pédagogique

Filière : aucune

Niveau : aucun

Autres descriptions : Langue : Français

Mention d'illustration

ill.

THE EDINBURGH STEREOSCOPIC ATLAS OF ANATOMY.

UPPER LIMB.

SURFACE ANATOMY—No. 3.

UPPER PART OF THE BACK.

The spines of the vertebrae lie at the bottom of a furrow between the erector spinae and other muscles on each side.

The spine of the vertebra prominens (seventh cervical) can usually be detected, but that of the first dorsal vertebra immediately below it is often equally distinct. The other dorsal spines can only be distinguished with difficulty.

The outline of the scapula is obscured by muscles covering it, but the lower angle can be made out by palpation, and the spine and acromion process can also be readily felt. About the scapula itself there are several hollows and elevations, which vary very considerably in different individuals on account of the varying sizes of different muscles.

The upper margin of the trapezius slopes outwards from the neck to the clavicle (4) and some of its fleshy fibres below this level form an elevation just above the spine of the scapula (1).

When the arms are in the attitude shown here, the scapula is rotated, so that the inner end of the spine is displaced downwards, and the lower angle of the bone rotated outwards. This accounts for the apparently low position of the spine of the scapula. Below the spine is seen a rounded elevation showing the position of the rhomboideus major muscle. The root of the spine of the scapula usually lies at the level of the second dorsal vertebra, and the lower angle of the scapula extends down to the seventh rib. The 'triangle of auscultation' lies to the inner side of this lower angle.

The figures indicate—

- | | | |
|------------------------|-------------------------------|------------------------------|
| 1. Trapezius muscle. | 3. Spine of scapula. | 6. Rhomboideus major muscle. |
| 2. Vertebra prominens. | 4. Upper margin of trapezius. | 7. Acromion process. |
| | 5. Deltoid muscle. | |

UPPER LIMB—SURFACE ANATOMY. No. 3.
EDINBURGH STEREOSCOPIC ANATOMY.
Copyright: T. C. & E. C. Jack, Edinburgh, & 34 Henrietta St., London, W.C.

