
192. The section is slightly to the right of the Mesial Plane in front. The Edinburgh Stereoscopic Atlas of Anatomy. Head and Neck. Sagittal section of the Head - N°1.

Numéro d'inventaire : 1979.26251.10

Type de document : photographie

Éditeur : Edinburgh Stereoscopic Anatomy, T.C. & E.C. Jack (34, Henrietta St., London, W.C. Edinburgh, London)

Période de création : 1er quart 20e siècle

Date de création : 1900 (vers)

Description : Epreuves stéréoscopiques positives gélatino-argentiques contrecollées sur un carton rigide contenant un texte descriptif (format du carton : 230 x 180). Série rangée dans un emboîtement en carton sous forme de reliure en deux parties avec la mention "Pestalozzi Stereographs. Anatomy" sur la tranche.

Mesures : hauteur : 90 mm ; largeur : 180 mm

Notes : Descriptif : coupe du crâne (anatomie).

Mots-clés : Méthodes pédagogiques actives (y compris la coopération scolaire, classes vertes, méthode Freinet)

Pratique pédagogique

Filière : aucune

Niveau : aucun

Autres descriptions : Langue : Français

Mention d'illustration

ill.

THE EDINBURGH STEREOSCOPIC ATLAS OF ANATOMY.

HEAD AND NECK.

SAGITTAL SECTION OF THE HEAD—No. 1.

THE SECTION IS SLIGHTLY TO THE RIGHT OF THE MESIAL PLANE IN FRONT, AND HAS LEFT THE SEPTUM OF THE NOSE, BUT AT THE BACK IT IS SLIGHTLY TO THE LEFT OF THE MESIAL PLANE.

The disc of cartilage between the sphenoid and the occipital bone is seen, as the subject was young, and it can also be noticed that the odontoid process of the axis vertebra is tipped with cartilage. Several of the large subarachnoid spaces at the base of the brain are seen (*vide* Spinal Canal, No. 2), the *cisterna basalis* a large cavity in front of the pons Varolii, the *cisterna pontis*, which is the continuation upwards of the anterior part of the subarachnoid space of the spinal canal, and the *cisterna magna*, between the under surface of the cerebellum and the back of the medulla. Where the section passes through the larynx, the true vocal cord is seen, with the laryngeal recess immediately above it, and the false vocal cord immediately above that again.

In the child the larynx is at a higher level than in the adult, so that the cricoid cartilage lies at the level of the fifth cervical vertebra, while in the adult it is usually at the sixth.

The adenoid tissue in the pharyngeal wall in front of the basi-occipital bone is somewhat hypertrophied in this specimen, forming the well known condition of adenoid vegetations. These vegetations spread outwards on each side to the region of the lateral recess of the pharynx.

The hyoid bone can be seen in section just in front of the figure 15. The interval between it and the symphysis menti corresponds to the floor of the mouth.

The figures indicate—

- | | | |
|-----------------------------|------------------------------|--------------------------------|
| 1. Nasal septum. | 7. Fourth cervical vertebra. | 13. Base of tongue. |
| 2. Basi-sphenoid. | 8. Superior maxilla. | 14. Lateral recess of pharynx. |
| 3. Basi-occipital. | 9. Mandible. | 15. Epiglottis. |
| 4. Atlas, anterior arch. | 10. Cisterna basalis. | 16. True vocal cord. |
| 5. Axis. | 11. Cisterna pontis. | 17. Cricoid cartilage. |
| 6. Third cervical vertebra. | 12. Cisterna magna. | |

HEAD AND NECK—SAGITTAL SECTION. No. 1.
(EDINBURGH STEREOSCOPIC ANATOMY.)
Copyright: T. C. & E. C. Jack, Edinburgh, & 34 Henrietta St., London, W.C.

